

Coronal X-ray emission from an intermediate-age brown dwarf

Yohko Tsuboi

*Department of Science and Engineering, Chuo University, Kasuga 1-13-27, Bunkyo-ku,
Tokyo 112-8551, Japan*

tsuboi@phys.chuo-u.ac.jp

Yoshitomo Maeda

*High Energy Astrophysics Division, Institute of Space and Astronautical Science,
Yoshinodai 3-1-1, Sagami-hara, Kanagawa 229-8510, Japan*

ymaeda@astro.isas.ac.jp

Eric D. Feigelson, Gordon P. Garmire, George Chartas, Koji Mori

*Department of Astronomy and Astrophysics, The Pennsylvania State University, 525
Davey Laboratory, University Park, PA 16802, USA*

edf@astro.psu.edu, garmire@astro.psu.edu, chartas@astro.psu.edu,
mori@astro.psu.edu

Steven H. Pravdo

*Jet Propulsion Laboratory, Mail Stop 306-438, California Institute of Technology,
Pasadena, CA 91109, USA*

spravdo@jpl.nasa.gov

ABSTRACT

We report the X-ray detection of the brown dwarf (BD) companion TWA 5B in a $\simeq 12$ Myr old pre-main sequence binary system. We clearly resolve the faint companion (35 photons) separated from the X-ray luminous primary by 2 arcsec in a *Chandra* ACIS image. TWA 5B shows a soft X-ray spectrum with a low plasma temperature of only 0.3 keV and a constant flux during the 3 hour observation, of which the characteristics are commonly seen in the solar corona. The X-ray luminosity is 4×10^{27} erg s $^{-1}$ (0.1–10 keV band) or $\log L_X/L_{bol} = -3.4$.

Comparing these properties to both younger and older BDs, we discuss the evolution of the X-ray emission in BDs. During their first few Myr, they exhibit

high levels of X-ray activity as seen in higher mass pre-main sequence stars. The level in TWA 5B is still high at $t \simeq 12$ Myr in $\log L_X/L_{bol}$ while kT has already substantially cooled.

Subject headings: stars: coronae — stars: individual (TWA 5B, CoD $-33^\circ 7795$ B) — stars: brown dwarf — X-rays: stars

1. Introduction

With masses below the hydrogen burning mass limit $M < 0.08 M_\odot = 80 M_{jup}$ and cool, nearly neutral surfaces, brown dwarfs (hereafter BDs) had been expected to lack high energy activity. However, X-rays from several young BDs with ages $t \sim 1$ Myr old were discovered with *ROSAT* in nearby star forming regions (Neuhäuser & Comerón 1998; Neuhäuser, Briceno & Comerón. 1999, Mokler & Stelzer 2002). After the launch of the *Chandra X-ray Observatory*, the number of X-ray-detected young BDs has increased an order of magnitude (Garmire et al. 2000, Imanishi, Koyama & Tsuboi 2001, Imanishi, Tsujimoto & Koyama 2001, Preibisch & Zinnecker 2001, Feigelson et al. 2002a). In young stellar clusters around 1 Myr old, hard and variable X-rays are now detected from about 1/4 of the BD population near the ‘saturation level’ of $\log L_X/L_{bol} \simeq -3$. These properties are indistinguishable from those of higher mass young objects which will evolve into dM stars.

Nearly all field L- and T-type older BDs with ages $t \simeq 500$ –1000 Myr lack $H\alpha$ emission, a classical signature of active regions and plage, but exhibit radio flares (Gizis et al. 2000, Berger 2002). The weak $H\alpha$ in field BDs is particularly surprising in light of their rapid rotation (Basri 2001). One field BD at age $t \simeq 500$ Myr, however, is clearly magnetically active. LP944-20 showed a 1–2 hr flare during a *Chandra* observation with a peak luminosity $L_X = 1 \times 10^{26}$ erg s $^{-1}$ ($\log L_X/L_{bol} \simeq -4.1$) with a surprisingly soft spectrum, while it exhibited no detectable quiescent X-rays before the flaring (Rutledge et al. 2000). This BD also has exceptionally strong radio emission (Berger et al. 2001) and shows $H\alpha$ emission of $EW \sim 1$ Å (Tinney 1998). At the intermediate age of $t \simeq 100$ Myr, three probable BDs in the Pleiades have been observed with *XMM – Newton* of which one, Roque 9, may have been detected at $\log L_X/L_{bol} \simeq -2.6$ (Briggs & Pye 2002, Stelzer & Neuhäuser 2002).

The evolution of magnetic activity in BDs thus appears confusing: $H\alpha$ and X-rays appear to decay as they descend the Hayashi tracks and cool, while radio emission does not. The nonthermal radio emission is particularly puzzling with both flare and persistent components far stronger than expected from the well-established Güdel-Benz radio/X-ray relationship that holds for all other late-type stars (Berger 2002). This contrasts with the

theoretical expectation that BD surface magnetic activity should decay with age due to increased resistivity of the atmosphere (Mohanty et al. 2002). The evolution of magnetic activity in higher mass late-type stars is also not fully understood. Various multiwavelength manifestations of magnetic activity (X-ray flares, radio flares, $H\alpha$ emission, large starspots, strong surface fields, etc.) are consistently seen at high levels in T Tauri stars, but do not appear to depend on rotation as seen in main sequence stars (Feigelson & Montmerle 1999, Feigelson et al. 2003).

We introduce to this issue the *Chandra* detection of TWA 5B, a BD at the intermediate age of $t \simeq 12$ Myr. TWA 5B is a faint red object ($H = 12$ mag) located $2''$ north of and co-moving with TWA 5A which itself is a spectroscopic binary (Lowrance et al. 1999, Webb et al. 1999, Neuhäuser et al. 2000). This system is part of the dispersed TW Hydra Association (TWA) of T Tauri stars, all of which are likely outliers of the giant Sco-Cen OB association (Kastner et al. 1997, Mamajek et al. 2000). TWA 5 is the first identified PMS star-BD binary system with an estimated age of 12 ± 6 Myr (Weintraub et al. 2000). We assume here the mean Hipparcos distance to the 4 brightest TWA members of 55 pc. Optical and near-infrared photometry of TWA 5B gives a spectral type of M8.5–M9 and bolometric luminosity $\log L_{bol}/L_{\odot} \simeq -2.6 \pm 0.3$, giving an inferred mass between 15 and 40 M_{jup} (Neuhäuser et al. 2000). The equivalent width of $H\alpha \sim 20 \text{ \AA}$ lies between those of young BDs (10–100 $\text{\AA}$ at ~ 1 Myr old) and those of field BDs ($\lesssim 1 \text{ \AA}$ at ~ 500 –5000 Myr). TWA 5B is thus clearly an intermediate-age BD and may be near the end of its brief deuterium burning phase (Weintraub et al. 2000). The inferred mass is not far from the boundary between a brown dwarf and a planet of about 12 M_{jup} .

2. Observations and results

TWA 5B was observed on 2001 April 15 with the *Chandra X-ray Observatory* for 10.3 ks using the ACIS-S detector (see Weisskopf et al. 2002 for a description of the satellite and instrument). The S3 CCD chip was operated in a rapid readout subarray mode to reduce photon pileup of the primary TWA 5A which was known from *ROSAT* studies to be very bright (Webb et al. 1999). The more frequent readouts resulted in a net on-target exposure of 9.3 ks. Image analysis started with Level 1 events from which we removed a $\pm 0''.25$ positional randomization, filtered with Good Time Intervals, and corrected charge transfer inefficiency problem as described by Townsley et al. (2000). To achieve the highest possible spatial resolution, we applied a correction to estimate the arrival location of each photon within each $0''.5$ pixel based on the precise distribution of charge received in adjacent pixels (Tsunemi et al. 2001, Mori et al. 2001). Spectral analysis was performed using Level 2 events

provided by the standard satellite pipeline processing.

Visual examination of the raw (Level 2) ACIS S3 CCD image shows very strong emission with about 6200 counts from TWA 5A, a CCD readout stripe oriented NE/SW, and a faint excess of events $2''$ to the north where TWA 5B lies. The TWA 5B emission becomes much clearer after the special processing steps outlined above. Figure 1 (center) shows this X-ray image in a very soft 0.1–1.2 keV band that highlights the emission from the BD companion. Note that a resolution of $\simeq 0''.3$ FWHM is achieved using the subpixel process, while the original data has $\simeq 0''.5$ FWHM (these FWHM values are measured from radial profile binned by 0.05 pixel.)

We extracted 35 photons from standard Level 2 event list around TWA 5B in the 0.1–8 keV band using a $0''.7$ radius circle after removal of background from an annulus around TWA 5A (see Figure 1, right panel). Since the extracted region is smaller than the standard *Chandra* point spread function at this position, a modified ‘auxiliary reference file’ was created. The background-subtracted spectrum of TWA 5B was fitted with an optically thin MEKAL thermal plasma model with elemental abundances fixed at 0.3 times of the solar value, based on the fitting results of other young stellar objects (YSOs) detected by *Chandra* and *ASCA* (e.g. Imanishi, Koyama, & Tsuboi 2001). The resulting spectrum, shown in Figure 2, is unusually soft with peak emission at 0.7 keV and no photons above 1.5 keV. Best-fit parameters with 90% confidence levels are: temperature $kT = 0.3^{+0.2}_{-0.1}$ keV, line-of-sight column density $N_H < 6 \times 10^{20} \text{ cm}^{-2}$, emission measure $4 \pm 1 \times 10^{50} \text{ cm}^{-3}$, and luminosity $L_X = 4 \times 10^{27} \text{ erg s}^{-1}$ in the 0.1–8 keV band. The spectral fit was essentially perfect with reduced chi-squared of 0.5 for 3 degrees of freedom. The plasma temperature and the luminosity are close to typical plasma conditions in the Sun today: at solar maximum, typical temperatures are $kT \simeq 0.2\text{--}0.5$ keV with $L_X \simeq 1 \times 10^{27} \text{ erg s}^{-1}$ (Peres et al. 2000). During a powerful solar flare, the typical temperatures rise to $T \simeq 1\text{--}2$ keV, much hotter than seen in TWA 5B but typical of other PMS stars and BDs (e.g. Feigelson et al. 2002a).

A light curve of the photon arrival times was examined and no significant variations were found during the 11 ks duration of the observation. We can exclude variations greater than 50% in amplitude over the 10 ks observation.

3. Discussion

3.1. Evolution of X-ray emission

As outlined in the Introduction, there is considerable confusion in our understanding of the evolution of BD magnetic activity. Figure 3 shows our current knowledge of the rela-

tionship between X-ray emission, parameterized as L_X/L_{bol} and the age of BDs. The X-ray properties for BDs in the Orion Nebula, ρ Ophiuchi and IC 348 are obtained from *Chandra* ACIS data by Feigelson et al. (2002a), Imanishi, Tsujimoto & Koyama (2001), and Preibisch & Zinnecker (2001), respectively. The field BD LP 944-20 detection is from Rutledge et al. (2000). Individual BD ages are distributed randomly for one order of magnitudes centered on 0.4 Myr (ρ Oph, Orion) and 1.5 Myr (IC348) based on Luhman et al. (2000). For Orion, the upper-limits of X-ray undetected brown dwarfs were obtained as described in Feigelson et al. (2002a, equations 7 and 10) assuming negligible absorbing column and bolometric luminosities from Hillenbrand and Carpenter (2000). The data of LP 944-20 are separately shown for the flare, the upper limit at the quiescent (pre-flare) phase and the time-average during full observation, while the others are shown only for the last one (full observations). Except for LP 944-20, only two sources in ρ Oph and one in IC348 have clear flares with amplitudes less than a factor of ten, which are within the range of the data scattering of each field. TWA 5B at $\log L_X/L_{bol} = -3.4$ lies just below the -3.0 saturation limit in main sequence clusters and is consistent with the BDs seen in younger clusters.

Although a clear conclusion cannot be made from one object, this result is consistent with a continued high level of BD X-ray surface activity through $t \simeq 12$ Myr. The decay to lower levels would then occur during the 10^7 – 10^8 yr phase, considering the averaged level of LP 944-20. This is consistent with the nearly constant (or even slightly rising; see Feigelson et al. 2003) $\log L_X/L_{bol}$ distribution from $t < 1$ Myr through $t \simeq 20$ Myr seen in higher mass late-type stars. The high level in $\log L_X/L_{bol}$ in younger phase might be related to the deuterium burning, with the fact that TWA 5B lies just at the end of the burning phase (Weintraub et al. 2000).

A somewhat different result emerges from consideration of the relationship between X-ray plasma temperature and age in BDs (Figure 4). Here the temperatures are derived from spectral fits to *Chandra* ACIS data by the authors listed above. While very young BDs have hotter temperatures above 1 keV, the intermediate-age BD TWA 5B is significantly cooler at $kT \simeq 0.3$ keV similar to the temperature found in the flare of the older LP 944-20 (Rutledge et al. 2000). However, the low temperature in TWA 5B, combined with the lack of variability, suggests that its X-ray emission arises more from a ‘corona’ than a single ‘flare’.

While a similar trend from hotter plasma to cooler plasma is seen in higher-mass YSOs, the timescale for this transition is much longer than implied for BDs. X-ray temperatures are typically $kT \simeq 6$ keV for protostars (e.g. Imanishi, Koyama, & Tsuboi 2001) and $kT \simeq 1 - 4$ keV for typical T Tauri stars (e.g. Feigelson et al. 2002a), with highly variable flare-dominated lightcurves. This hotter plasma is dominated by cooler coronal plasma only after $\simeq 500$ Myrs (Güdel, Guinan & Skinner 1997).

It is interesting that the plasma temperature in LP 944-20 flaring is similar to that in TWA 5B quiescent. Considering that the X-ray luminosity of the LP 944-20 flaring is at least an order of magnitudes less than that of TWA 5B quiescent, the X-rays of TWA 5B, which appear like coronal emission, might be explained by a superposition of a number of such small flares. The similar possibility has been suggested and examined for the solar corona (Parker 1988, Shimizu & Tsuneta 1997).

3.2. Comments on the relationship between X-ray and H α emissions

Figure 5 shows our current knowledge of the relationship between X-ray emission, parameterized as L_X/L_{bol} and the H α equivalent widths of BDs. In this plot, the ordinate indicates the X-ray surface brightness at the BD surface, while the abscissa represents the H α surface brightness, in the zero's order. Here the H α properties are obtained from Jayawardhana, Mohanty & Basri (2002) for BDs in the ρ Ophiuchi, Luhman (1999) and Herbig (1998) for those in IC 348, and Tinney (1998) for the field BD LP 944-20. For the time-averaged X-rays of detected BDs, we obtained a correlation of $\log(L_X/L_{bol}) = 1.5 \times \log(H\alpha) - 5.3$. Such correlation has been already known in higher mass weak-lined T Tauri stars (for a recent example; Preibisch & Zinnecker 2002), which is interpreted by the hypothesis that the chromosphere (traced by the H α activity) is heated by a sufficient overlying corona (X-ray activity) (Cram 1982).

Figure 5 also shows that the strongest six (~ 100 Å) in the H α equivalent width were not detected in X-rays. As with higher mass classical T Tauri stars, these strong emission lines would be accompanied by IR disks. The relation given in Figure 5 is thus straightforwardly interpreted by the scenario that X-rays are mainly due to the coronal activity while H α is dominated by the chromospheric activity ($\lesssim$ a few tens Å) and by the disk activity ($\gtrsim$ a few tens Å). These nondetections also suggest that BD X-ray activity does not likely originate from star-disk interaction but from the solar-like magnetic activity.

4. Conclusion

We report the detection of X-rays from an intermediate-age ($t \simeq 12$ Myr) BD at a level of $L_X \simeq 4 \times 10^{27}$ erg s $^{-1}$ (0.1–10 keV band) or $\log L_X/L_{bol} = -3.4$. The X-ray spectrum is very soft; the dominant plasma temperature is only 0.3 keV. No variability is seen during the 10.3 ks observation. Those characteristics are common to those in the solar corona. Our observation provides a link between the active state seen in younger $t \simeq 1$ Myr stellar

clusters, and a relatively inactive state, showing continuity in the evolution of both X-ray luminosity and plasma temperature: $\log L_X/L_{bol}$ has not yet decayed by $t \simeq 12$ Myr while kT has already substantially cooled. The correlation between X-rays and $H\alpha$ implies the idea that the X-rays are mainly due to the coronal activity while $H\alpha$ is dominated by the chromospheric activity ($\lesssim$ a few tens Å) and by the disk activity ($\gtrsim$ a few tens Å). Finally, we note that since TWA 5B is not far from the boundary between BDs and the most massive planets found orbiting nearby stars, it raises the possibility that massive planets might emit X-ray during their youth.

We express our thanks to all the *Chandra* team for many efforts of the fabrication of satellite, launching, daily operation, software developments and calibrations. We also thank Leisa Townsley for supplying the CTI corrector of ACIS and Kensuke Imanishi the referee Gibor Basri for useful comments on brown dwarf flares and coronae. This research was supported by NASA contract NAS 8-38252.

REFERENCES

- Basri, G. 2001, in ASP Conf. Ser. 223: Cool Stars, Stellar Systems and the Sun, 261 (Ramon J. Garcia Lopez, Rafael Rebolo, and Maria Rosa Zapaterio Osorio. eds),
- Berger, E., Ball, S., Becker, K. M., Clarke, M., Frail, D. A., Fukuda, T. A., Hoffman, I. M., Mellon, R., Momjian, E., Murphy, N. W., Teng, S. H., Woodruff, T., Zauderer, B. A., Zavala, R. T. 2001, *Nature*, 410, 338
- Berger, E. 2002, *ApJ*, 572, 503
- Briggs, K. R. & Pye, J. P. 2002, in *Stellar Coronae in the Chandra and XMM-Newton Era* (F. Favata & J. Drake, eds), ASP Conf. Ser., in press (astro-ph/0110109)
- Feigelson, E. D. & Montmerle, T. 1999, *ARA&A*, 37, 363
- Feigelson, E. D., Broos, P., Gaffney, J. A., Garmire, G., Hillenbrand, L. A., Pravdo, S. H., Townsley, L., Tsuboi, Y. 2002a, *ApJ*, 574, 258
- Feigelson, E. D., Gaffney, J. A., Garmire, G., Hillenbrand, L. A. & Townsley, L. 2003, to appear in *ApJ*(astro-ph/0211049)
- Garmire, G., Feigelson, E. D., Broos, P., Hillenbrand, L. A., Pravdo, S. H., Townsley, L., Tsuboi, Y. 2000, *AJ*, 120, 1426
- Gizis, J. E., Monet, D. G., Reid, I. N., Kirkpatrick, J. D., Liebert, J., & Williams, R. J. 2000, *AJ*, 120, 1085
- Güdel, M., Guinan, E. F., & Skinner, S. L. 1997, *ApJ*, 483, 947
- Herbig, G. H. 1998, *ApJ*, 497, 736
- Hillenbrand, L. A. & Carpenter, J. M. 2000, *ApJ*, 540, 236
- Imanishi. K., Koyama, K., Tsuboi, T. 2001, *ApJ*, 557, 747
- Imanishi. K., Tsujimoto, M., Koyama, K. 2001, *ApJ*, 563, 361
- Jayawardhana, R., Mohanty, S., & Basri, G. 2002, *ApJ*, 578, L141
- Kamata, Y., Koyama, K., Tsuboi, Y., & Yamauchi, S. 1997, *PASJ*, 49, 461
- Kastner J. H., Zuckerman, B., Weintraub, D. A., Forveille, T. 1997, *Science*, 277, 67
- Lowrance, P. J., et al. 1999 *ApJ*, 512, L69

- Luhman, K. L. 1999, *ApJ*, 525, 466
- Luhman, K. L., Rieke, G. H., Young, E. T., Cotera, A. S., Chen, H., Rieke, M. J., Schneider, G., Thompson, R. I. 2000, *ApJ*, 540, 1016
- Mamajek, E. E., Lawson, W. A., & Feigelson, E. D. 2000, *ApJ*, 544, 356
- Mohanty, S., Basri, G., Shu, F., Allard, F., & Chabrier, G. 2002, *ApJ*, 571, 469
- Mokler, F. & Stelzer, B. 2002, *A&A*, 391, 1025
- Mori, K., Tsunemi, H., Miyata, E., Baluta, C. J., Burrows, D. N., Garmire, G. P., & Chartas, G. 2001, in *ASP Conf. Ser. 251: New Century of X-ray Astronomy*, 576
- Neuhäuser, R. & Comerón, F. 1998, *Science*, 282, 83
- Neuhäuser, R., Briceno, C., & Comerón, F., 1999, *A&A*, 343, 883
- Neuhauser, R., Guenther, E. W., Petr, M. G., Brandner, W., Huélamo, N., Alves, J. 2000, *A&A*, 360, L39
- Peres, G., Orlando, S., Reale, F., Rosner, R., & Hudson, H. 2000, *ApJ*, 528, 537
- Preibisch, T. & Zinnecker, H. 2001 *AJ*, 122, 866
- Rutledge, R. E., Basri, G., Martin, E. L., Bildsten, L. 2000, *ApJ*, 538, L141
- Simizu, T. & Tsuneta, S. 1997, *ApJ*, 486, 1045
- Stelzer, B. & Neuhauser, R. 2002, in *Brown Dwarfs*, IAU Symposium 211, in press (astro-ph/0206284)
- Tinney, C. G. 1998, *MNRAS*, 296, L42
- Townsley, L. K., Broos, P. S., Garmire, G. P., & Nousek, J. A. 2000, *ApJ*, 534, L139
- Tsunemi, H., Mori, K., Miyata, E., Baluta, C. J., Burrows, D. N., Garmire, G. P., & Chartas, G. 2001, *ApJ*, 554, 496
- Webb, R. A., Zuckerman, B., Platais, I., Patience, J., White, R. J., Schwartz, M. J., & McCarthy, C. 1999, *ApJ*, 512, L63
- Weisskopf, M. C., Brinkman, B., Canizares, C., Garmire, G., Murray, S., & Van Speybroeck, L. P. 2002, *PASP*, 114, 1

Weintraub, D. A., Saumon, D., Kastner, J. H., & Forveille, T. 2000, ApJ, 530, 867




Fig. 1.— The TWA 5 system showing the intermediate-age BD TWA 5B just north of the primary star: (left) I-band image from Neuhäuser et al. 2000; (center) *Chandra* x-ray image (0.1–1.2 keV) obtained here; and (right) diagram of the X-ray spectral extraction regions. The *Chandra* image is displayed in $0''.1$ pixels after special processing steps outlined in the text and smoothing with a $0''.5$ FWHM Gaussian.




Fig. 2.— The ACIS spectrum of TWA 5B extracted from a $0''.7$ radius region after background subtraction (see Figure 1, right panel). The spectrum is fitted with a coronal plasma model, and the lower panel shows the residuals from the best-fit model.




Fig. 3.— $\log L_X / L_{bol}$ vs. age of BDs. Symbols: Orion Nebula Cluster (open squares), ρ Ophiuchi (open triangles), and IC 348 (filled squares). The star indicates TWA 5B. LP 944-20 is separately shown for each period of the flare (the dotted circle), quiescent (the open circle), and full observation (the arrow). Nondetections are indicated with arrows. All data are obtained from *Chandra* ACIS observations (see text for references).




Fig. 4.— Plasma temperature vs. age of BDs. The symbols are the same as those in Figure 3.

